

# NUR 2474 PHARMACOLOGY FOR PROFESSIONAL NURSING

## Pharm Mod 8 Quiz

The healthcare provider orders the following:

Prednisone 5 mg PO q6h × 4 days

5 mg PO q12h × 3 days

5 mg PO qDay × 3 days

Dispense 5-mg tablets

After completing 6 days of treatment the patient calls and asks how many pills should be left in the bottle because some may have been lost. How many pills should be left?

- A. 4
- B. 6
- C. 10
- D. 5**

learnexams

A nurse obtains a health history from a patient who has gout and is taking a glucocorticoid. The nurse should follow up on which finding?

- A. Blood glucose level of 140 mg/dL**
- B. Flushing and urticaria
- C. 6-kg weight loss
- D. Heart rate of 88 beats/min

After administering acetylcysteine [Mucomyst] to a patient who overdosed on acetaminophen [Tylenol], a nurse should recognize which outcome as an indicator of the therapeutic effects of acetylcysteine?

- A. Absence of jaundice**
- B. Clear breath sounds
- C. Increased bowel sounds
- D. Palpable pedal pulses

When evaluating the effects of probenecid, the nurse should monitor which laboratory result?

- A. Uric acid level**
- B. Blood pH
- C. Sodium level
- D. Hemoglobin

learnexams

A patient develops hypotension, laryngeal edema, and bronchospasm after eating peanuts. Which medication should the nurse prepare to administer?

- A. Hydroxyzine [Vistaril]
- B. Epinephrine**
- C. Promethazine [Phenergan]
- D. Diphenhydramine [Benadryl]

The nurse identifies which symptom as a result of activation of histamine<sub>1</sub> (H<sub>1</sub>) by allergic reaction?

- A. Sweating
- B. Pupillary dilation
- C. Bronchoconstriction**
- D. Hypertension