NSG 3100 EXAM 3 LATEST 2023-2024 (2 VERSIONS A & B) ACTUAL EXAM ALL 180 QUESTIONS AND CORRECT DETAILED ANSWERS WITH RATIONALES|ALREADY GRADED A+

VERSION A

When changing the colostomy appliance, cut the opening in the skin barrier no more than ____ larger than the stoma. - ANSWER- 1/8 inch

The best time to change a pouching system is in the morning or 2 to 4 hours after meals. True or False - ANSWER- True

An unlicensed assistive personnel (UAP) reports to the nurse that a client being fed experienced coughing and choking when swallowing. The client states, "It feels like the food is stuck in my throat." What does the nurse suspect is happening with this client? - ANSWER- The client is having dysphagia.

While undergoing a soapsuds enema, the client complains of mild abdominal cramping. The nurse should: - ANSWER- lower the bag

The client has an indwelling catheter. The nurse should obtain a sterile urine specimen by: - ANSWER- syringe to withdraw from cath port

limiting fluids has what effect on urine? - ANSWER- raises specific gravity

The nurse is alert to the possibility that for 24 to 48 hours after the postoperative procedure, clients may experience the following as a result of the anesthetic used during the surgery: - ANSWER- paralytic ilyus

before fecal occult test eat - ANSWER-bread

diagnosis of Alteration in urinary elimination, retention. On assessment, the nurse anticipates that this client will exhibit: - ANSWER- a feeling of pressure and voiding of small amounts.

diarrhea patients should consume - ANSWER- lean meats

learnexams

A colonoscopy is ordered and the patient has questions about the examination. Before the colonoscopy, the nurse teaches the patient that:
- ANSWER- light sedation is normally used

In an assessment of a client with overflow incontinence, the nurse expects to find that the client has: - ANSWER- constant dribbling of urine

A nurse is monitoring a client's fluid intake. For breakfast, the client consumed 8 oz of milk, 10 oz of water, 4 oz of flavored gelatin, 1 scrambled egg, 1 crisp piece of bacon, and 2 biscuits with jelly. The nurse should record the fluid intake as: - ANSWER- 660 mLs

24 hour urine needs to be restarted in the event that: - ANSWER- client voids in the toilet

Which interventions prevent or minimize the risk factors in clients at risk for spiration? - ANSWER- Feeding the patient small bites.

Keeping suction equipment nearby.

Positioning the client upright at 90 degrees.

Providing liquids with a thickening agent.

The client receiving a tube feeding develops diarrhea. The nurse should: - ANSWER- adjust rate of infusion

Critical care element for clients with ileostemy - ANSWER- skin care

Why would a nurse order a guaiac test - ANSWER-blood in stool

most important question for MRI - ANSWER- Any metal in body?

considerations for diagnostic procedures - ANSWER- what type of sample?

NPO?

Contrast?

Medications?

Duration of test

consent?

what is included in the CBC - ANSWER- Hgb

Hct

Erythrocytes

RBC

Leukocytes

diff wbc

Components of BMP (basic metabolic panel) - ANSWER-BUN

Creatinine

Potassium

Sodium

learnexams

Calcium

Chloride

Glucose

Components of a CMP (Comprehensive Metabolic Panel) - ANSWER-Liver

Livei

Proteins

albumin

bilirubin