

**Phtls Pre and Post Test Answers Update
2023-2024 New Latest Version Best
Studying Material Actual Past Exam**

Medical Surgical Nursing 1 (Santa Fe College)



[shutterstock.com](https://www.shutterstock.com) · 16/539132/14

PHTLS Pre & Post Test Questions & Answers_ All correct.

The displacement of tissue away from the path of a projectile, both temporarily and permanently, is known as:

- A. Conization
- B. Cavitation**
- C. Crepitation
- D. Contusion

The single most important factor in determining the potential for injury due to energy exchange is:

- A. Mass of the bodies involved
- B. Velocity of the bodies involved**
- C. Density of the tissues involved
- D. Surface area of the impact involved

In the management of shock, isotonic crystalloid solutions, such as Ringer's, are preferred because:

- A. The protein molecules in crystalloid solutions act as volume expanders
- B. These fluids draw interstitial fluid into the vascular space to enhance volume
- C. These solutions will stay in the vascular space longer than water solutions, such as D5W**
- D. Their pH enhance oxygen delivery to the tissues

With respect to the distance of a fall, which of the following is a guideline for determining a critical fall?

- A. 3 times the height of the patient**
- B. 2 times the height of the patient

PHTLS Pre & Post Test Questions & Answers_ All correct.

- C. 5 times the height of the patient
- D. 1 ½ times the height of the patient

The phase of an explosion, or blast, in which hollow organs are squeezed and may rupture is called the _____ phase.

- A. Tertiary phase
- B. Quaternary phase
- C. Secondary phase
- D. Primary phase

During the primary survey and management of a trauma patient, the E in ABCDE stands for _____?

- A. Edema
- B. Eyes & ears
- C. Expose/Environment
- D. Electrical therapy

The time in which surgical intervention can make a difference in patient outcome is the _____?

- A. Golden period
- B. Golden time
- C. Golden era
- D. Golden minutes

In the absence of extenuating circumstances, the maximum amount of time it should take to identify and manage immediate threats to life, prepare the patient for transport and begin transport is _____?

- A. 5 minutes
- B. 10 minutes

PHTLS Pre & Post Test Questions & Answers_ All correct.